

Measurement of Transverse Single Spin Asymmetry A_N in Eta Mass Region at Large Feynman X_F with the STAR Forward Pion Detector

Steven Heppelmann, for the STAR Collaboration
Davey Lab, Penn State University, University Park, Pa. 16802

Abstract

The large values of the transverse single spin asymmetry, A_N , seen in forward π^0 production from polarized proton collisions have stimulated important questions and have been studied in QCD based transverse spin models. We report the first RHIC measurement of A_N for the two photon mass region of the η meson. The η peak for meson energies greater than 50 GeV (Feynman $x_F > 0.5$) was observed in the STAR Forward Pion Detectors, along with π^0 mesons, at pseudo-rapidity of 3.65 in $\sqrt{s} = 200\text{GeV}$ pp collisions. The π^0 transverse asymmetry, which has already been reported in detail by the STAR collaboration, is compared to the transverse asymmetry in the η mass region. The surprising observation is that in this kinematic region, the η asymmetry is larger than the already large π^0 asymmetry.

Key words: Transverse Single Spin Asymmetry, Eta Meson
PACS: 13.85.-t, 13.88.+e, 14.4.Aq

1. Introduction

Measurements of large transverse single spin asymmetries of high energy forward meson production in collisions between polarized protons have surprised physicists for many years. It is well known that such asymmetries require both a non-vanishing helicity flip amplitude and interference between real and imaginary amplitudes. From a perturbative QCD perspective, this means that these asymmetries do not occur in leading twist QCD with collinear factorization. It has been determined, however, that the π^0 spin averaged cross section is in fairly good agreement with Next to Leading Order Perturbative QCD[1]. Thus these transverse asymmetries provide a laboratory for the sensitive study of non-leading twist aspects of hadron interactions in processes where the underlying leading twist calculations have been shown to be meaningful.

Two basic parton model pictures have developed to explain how these asymmetries can come about. In the Sivers model[2,3], the initial transverse momentum distribution of large x quarks, in conjunction with absorptive effects, could lead to the required spin dependent transverse momentum that can explain these asymmetries. In the Collins model[4], corresponding higher twist effects are associated with the final state fragmentation process that produces the final state meson from a parton. The parton retains its initial state polarization through the hard scattering process.

Here we will consider a comparison between the forward asymmetries, A_N , measured with forward π^0 and η mesons. The central assumption of factorization, implicit in either the Collins or Sivers approach, leads to the conclusion that the hard scattering process determines the transverse momentum of the jet, independent of fragmentation details. However, jets of different types may favor fragmentation into different mesons. For example, it is widely accepted that if a u quark fragments with large z , it tends to fragment into a π^+ meson and a d quarks similarly tends to fragment into a π^- meson. It is believed that the transverse spin of the quarks (like the longitudinal spin) in a polarized proton is highly correlated with isospin of those quarks. The well known difference in sign of A_N for π^+ vs. π^- mesons may be evidence of this. When we flip the z component of the isospin (I_z) of the observed meson, we also tend to flip the transverse asymmetry A_N and apparently the isospin of the jet parton. If the pion I_z is highly correlated with A_N , it is natural to ask how, for the $I_z = 0$ mesons, does the asymmetry depend upon the total isospin I . Does the transverse single spin asymmetry differ for the $I = 0$ η meson and the $I = 1$ π^0 meson? Of course the η and π^0 differ not only in isospin but also in the strangeness content of the wave function,

$$\pi^0 = \frac{1}{\sqrt{2}}(u\bar{u} - d\bar{d}) \quad \eta \simeq \frac{1}{\sqrt{3}}(u\bar{u} + d\bar{d} - s\bar{s}) \quad (1)$$

where to obtain the simple expression for η , the η, η' mixing angle has been approximated as $\theta_P \simeq -19.4^\circ$ [5]. While QCD based models for transverse single spin asymmetries have been widely studied, these models have not yet been extended to make predictions for differences in A_N for η vs. π^0 production.

Observed differences in A_N for π^0 and η pseudo-scalar mesons have already been reported by E704 [6]. For Feynman x_F above 0.4, they reported a substantially larger asymmetry for η than π^0 mesons but the significance of the difference was marginal.

2. Measurement

With the STAR Forward Pion Detector (FPD), during the 2006 RHIC run with transversely polarized protons of energy $\sqrt{s} = 200 \text{ GeV}$, the transverse single spin asymmetry for π^0 production at large Feynman x_F has already been reported[7]. Using the same FPD apparatus and same data set, the corresponding A_N asymmetry is now shown for photon pairs in the η meson region. The selection of events is chosen to optimize FPD acceptance and yield for η mesons relative to π^0 s, cutting on two photon pseudo-rapidity (Y) and azimuthal angle (ϕ), $((Y - 3.65)^2 + \tan(\phi)^2) < (0.15)^2$. This cut corresponds to the centers of the two FPD calorimeters located downstream, both left and right, of the yellow polarized RHIC beam. The transverse polarization axis is oriented at $\phi = \pm \frac{\pi}{2}$ with an average polarization for the yellow beam of 56%. The FPD acceptance for two

photon decays improves for larger x_F in the η mass region and is significant for x_F above about 0.4. In Figure 1, the three pairs of mass plots correspond to a further selection of events with photon pair energies in the indicated ranges. The mass regions that will be associated with π^0 and η mesons are indicated with the vertical bands. It is for the events in these two mass bands that the single spin asymmetry is calculated as a function of x_F using the cross ratio method[7]. These asymmetries are shown in Figure 2.




Fig. 1. Two photon mass distributions are shown in 3 energy ranges with π^0 and η mass bands indicated. Note the log scale on the upper plots and that the linear scale on the lower plots emphasizes the η peak.




Fig. 2. The dependence of A_N on x_F is plotted for the π^0 and η mass bands defined by $m_{\pi^0} = 0.135 \pm 0.05$ GeV for π^0 and $m_\eta = 0.55 \pm 0.07$ GeV for η .

3. Results and Summary

Like the E704 measurement, the asymmetry A_N for the η meson mass region is larger than that for the π^0 region. The weighted average of this asymmetry over Feynman x_F in the range $0.55 < x_F < .75$ is $\langle A_N \rangle_\eta = 0.361 \pm 0.064$. In comparison, for the π^0 mass region the corresponding asymmetry is $\langle A_N \rangle_{\pi^0} = 0.078 \pm 0.018$. The errors here are statistical, with preliminary estimates of systematic errors much smaller. This difference in A_N between the η mass region and the π^0 region is more than four standard deviations. In the η region, we further note the trend to higher asymmetry at larger x_F , raising the question as to whether the asymmetry is approaching a maximal value of 1 in the high x_F end of this range. This result is consistent with the E704 measurement but is more significant with largest A_N at higher x_F than previously measured.

References

- [1] J. Adams et al. *Phys. Rev. Lett.*, 97:152302, 2006.
- [2] D. W. Sivers. *Phys. Rev.*, D43:261–263, 1991.
- [3] U. D'Alesio and F. Murgia. *Phys. Rev.*, D70:074009, 2004.
- [4] J. C. Collins, S. F. Heppelmann, and G. A. Ladinsky. *Nucl. Phys.*, B420:565–582, 1994.
- [5] A. Bramon, R. Escribano, and M. D. Scadron. *Eur. Phys. J.*, C7:271–278, 1999.

- [6] D. L. Adams et al. *Nucl. Phys.*, B510:3–11, 1998.
- [7] B. I. Abelev et al. *Phys. Rev. Lett.*, 101:222001, 2008.